

1 Arrival of extremely volatile-rich high-Mg magmas
2 changes explosivity of Mount Etna

3 **Vadim S. Kamenetsky***

4 *ARC Centre of Excellence in Ore Deposits and School of Earth Sciences, University of*
5 *Tasmania, Hobart, Tasmania 7001, Australia*

6 **Massimo Pompilio**

7 *Istituto Nazionale di Geofisica e Vulcanologia - Sezione di Pisa, Via della Faggiola, 32*
8 *Pisa, 56126 Italy*

9 **Nicole Métrich**

10 *Laboratoire Pierre Sue, CNRS-CEA, CE-Saclay, 91191 Gif sur Yvette, France*

11 **Alexander V. Sobolev**

12 *Max-Planck-Institut für Chemie, Postfach 3060, 55020 Mainz, Germany and Vernadsky*
13 *Institute of Geochemistry, Moscow 117975, Russia*

14 **Dmitry V. Kuzmin**

15 *Max-Planck-Institut für Chemie, Postfach 3060, 55020 Mainz, Germany and Institute of*
16 *Geology and Mineralogy SB RAS, Novosibirsk 630090, Russia*

17 **Rainer Thomas**

18 *GeoForschungsZentrum Potsdam, Telegrafenberg B 120, D-14473, Potsdam, Germany*

19 *E-mail: Dima.Kamenetsky@utas.edu.au.

20 **ABSTRACT**

21 The volcanic hazard potential of Mount Etna volcano is presently nourished by
22 long-lasting, powerful eruptions of basaltic magmas coupled with increased seismicity

and ground deformation, and the world's largest discharge of volcanic gases. The current evolutionary cycle of Mount Etna activity is consistent with subduction-related chemical modifications of the mantle source. Arrival of a new mantle-derived magma batch beneath the volcano has been hypothesized, but is still elusive among the erupted products. Here we demonstrate petrological and geochemical affinities between the magmas supplying modern eruptions and high-Mg, fall-stratified ("FS") basalts ejected violently ~4 Kyr ago. The "FS" primitive magmas (~13 wt% MgO) are characteristically volatile-enriched (at least 3.8 wt% H₂O and 3300 ppm CO₂), and bear trace element signature of a garnet-bearing, metasomatised source (high Gd/Yb, K/La, U/Nb, Pb/Ce, Ca/Al). They started crystallizing olivine (Fo₉₁), clinopyroxene (Mg# 92.5) and Cr-spinel deep in the plumbing system (> 5 kb), contributing to the cumulate piles at depth and to differentiated alkaline basalt and trachybasalt magmas in the shallow conduit. Continuous influx of mantle-derived, volatile-rich magmas, such as those that supplied the "FS" fallout, provides a good explanation for major compositional and eruptive features of Mount Etna.

Keywords: Etna, volcano, picrite, melt inclusions, volatiles, degassing, mantle, metasomatism.

INTRODUCTION

Mount Etna, positioned at the destructive margin marking the complex collision zone between the African and European plates (Gueguen et al., 1998), is the largest volcano in Europe and one of the most active in the world. The present edifice of Mt. Etna was constructed with fractionation products of mantle-derived magmas, starting ~0.5 Ma ago with eruption of tholeiitic lavas, followed by transitional lavas, eventually

trending to SiO₂-undersaturated, alkaline basalts with distinct trace element and isotope signatures (Armienti et al., 2004; Corsaro and Pompilio, 2004; Tanguy et al., 1997). The compositional changes of the Mt. Etna magmatic system were temporally and genetically linked to continuous modifications in the mantle/lithospheric source(s), caused by metasomatic enrichment in volatiles and incompatible trace elements, and melting-related depletion in basaltic components (Kamenetsky and Clocchiatti, 1996). Dramatic changes in both the source composition and the volcano's eruptive dynamics have been tentatively ascribed to modification of the adjacent subduction system (Schiano et al., 2001; Tonarini et al., 2001), driven by vertical asthenospheric flow in response to the roll-back of the subducting Ionian slab (Doglioni et al., 2001; Gvirtzman and Nur, 1999).

From a historical perspective, Mt. Etna shows increasing activity in recent times, with the frequency of explosive (doubled since the "Fall Stratified" (FS) eruption of high-magnesian magmas ~4 Kyr ago) and effusive eruptions increasing significantly (Coltelli et al., 2000). More recently, since the early 1970s, the magmatic output of Mt. Etna has increased by a factor of four, and more than 120 eruptions have been recorded in the past 20 yr (Branca and Del Carlo, 2005). Notably, the erupted magmas have become more potassic and richer in ⁸⁷Sr and ¹¹B (Armienti et al., 2004; Clocchiatti et al., 1988; Tanguy et al., 1997; Tonarini et al., 2001), and have been accompanied by enormous output of volcanic gases CO₂, SO₂ and halogens (Allard, 1997; Allard et al., 1991; D'Alessandro et al., 1997). Although, the present day magmas are increasingly magnesian and volatile-rich (Clocchiatti et al., 2004; Corsaro and Pompilio, 2004; Métrich et al., 2004; Spilliaert et al., 2006), and their radiogenic and stable isotope compositions require a mantle origin (Allard et al., 1991; Armienti et al., 2004; D'Alessandro et al., 1997; Marty et al., 1994;

Tonarini et al., 2001), their MgO (< 8.2 wt%) and Ni (< 50 ppm) contents, and forsterite contents of olivine phenocrysts (< 84 mol% Fo) are too low to have been in equilibrium with a typical peridotite mantle.

On the other hand, the indications of a primitive melt entering the plumbing system of Mt. Etna from a mantle reservoir are explicit. In addition to the isotope systematics, these include records of large piles of mafic cumulates extending to depths of 18 km beneath the volcano (De Gori et al., 2005) and the “excessive” (compared to amounts degassed from the erupted magmas) supply of volatiles to the volcanic clouds (Allard et al., 1991; D’Alessandro et al., 1997). As primitive magmas have rarely been ejected in the eruptive history of Mt. Etna (Corsaro and Pompilio, 2004) and have not reached the surface since the “FS” eruption, we used the latter to elucidate controls on recent and current volcanic activity.

“FS” ERUPTION: ROCKS, MINERALS AND MELT INCLUSIONS

The 3930 ± 60 BP (^{14}C age; Coltelli et al., 2000) “FS” eruption produced a pyroclastic fallout deposit of 0.183 km^3 covering the entire eastern flanks of Mt. Etna (Coltelli et al., 2005). The deposit reaches 110 cm in thickness at ~7 km from the summit craters, and its thickness and dispersal indicate that the eruption column reached significant heights (18–20 km (Coltelli et al., 2005)), and thus had high explosive (subplinian) energy. The erupted products are a highly vesiculated (30–60 vol%) scoria lapilli containing euhedral crystals of olivine (< 8%), clinopyroxene (< 3%) and Cr-spinel (< 0.1%). Despite relatively low amounts of ferromagnesian phenocrysts the rocks are high-Mg basalt to picrite in composition (12–17 wt% MgO; Table DR1 of the GSA Data Repository¹) with high Ni and Cr abundances (135–285 and 920–1330 ppm,

respectively), characteristic of primitive, unfractionated magmas. The rocks are extremely enriched in CaO relatively to Al_2O_3 ($\text{CaO}/\text{Al}_2\text{O}_3 = 1.1\text{--}1.2$) despite the low modal abundances of clinopyroxene phenocrysts. The “FS” rocks are depleted in heavy rare earth elements (HREE) and high-field strength elements (HFSE) relative to REE of higher and similar incompatibility, respectively (Fig. 1). They are also relatively enriched in large-ion lithophile elements (LILE – K and Sr) and Pb (Fig. 1). Depletions and enrichments demonstrated by the ratios of similarly incompatible elements (e.g., K/La, Nb/U, Ce/Pb, Ti/Gd) are typical of subduction-related magmas and commonly attributed to selective addition of $\text{LILE} > \text{LREE} > \text{HFSE}$ to a mantle peridotite by melts and/or fluids derived from a subducting crust. High Gd/Yb in the magmas can be taken as evidence of residual garnet in the source.

High-Mg compositions of the “FS” olivine phenocrysts (Fo 90–91 mol% in the majority of crystals) and clinopyroxene (Mg# 96–92.5) also point to the primitive nature of their parental magmas (see GSA Data Repository). Early crystallization of “FS” clinopyroxene, confirmed by the occurrence of clinopyroxene inclusions in most forsteritic olivine may reflect high Ca abundance in the parental magma and high-pressure crystallization conditions (Grove et al., 1992). Spinel microphenocrysts and inclusions in olivine have extremely high-Cr (Cr# 70–80 mol%) and moderately low-Ti (0.6–1.0 wt% TiO_2) compositions that are also typical of many primitive alkaline island arc magmas, including those from nearby Vulcano, Aeolian arc (Kamenetsky and Clocchiatti, 1996; Kamenetsky et al., 2001).

Olivine-hosted melt inclusions, analyzed for major, trace and volatile elements, using different microbeam techniques, and subsequently corrected for post-entrapment

modifications (see GSA Data Repository for the details of analytical methods and calculations) provide more insight into the nature of the “FS” parental magmas. Melt inclusions (up to 250 μm in size) have negative-crystal shape with scalloped edges and appear as brownish glass with a spherical vapor bubble (Fig. 3), which is sometimes larger than typical shrinkage bubbles (i.e., $> 5 \text{ vol\%}$). The bubble walls are invariably lined with tiny Ca-Mg carbonate crystals (Fig. 3).

Melt inclusion compositions (after correction for minor post-entrapment modifications; see GSA Data Repository) show a limited range in MgO ($12.6 \pm 0.3 \text{ wt\%}$) and match the bulk rock major and trace element compositions (Fig. 1, 2). Thus, the melt inclusions support our conclusions about high-Mg and high-Ca compositions of the “FS” parental melt with the trace element systematics reminiscent of subduction-related magmatism. Although the average composition of melt inclusions is indistinguishable from the average bulk rock composition, the observed chemical variability among individual inclusions, especially in trace elements (Fig. 1, 2), is very large and cannot be induced by crystallization as the compositions of host olivine grains are similar.

Melt inclusions have moderately high Cl and S (average 0.15 and 0.17 wt%, respectively), but are exceptionally enriched in H_2O and CO_2 for basaltic melts, in general, and primitive, high-Mg basalt, in particular (Fig. 3). The original CO_2 content of trapped melts was undoubtedly higher than measured in glasses, because carbon present in the bubbles (gaseous phase and carbonates) is not quantified. Importantly, H_2O does not correlate with elements of similar incompatibility (see large variability in trace elements in Fig. 2), and thus the effects of degassing cannot be excluded. Nevertheless,

the highest measured abundances of dissolved volatiles indicate very deep crystallization (at least 400–500 MPa, Fig. 3) of clinopyroxene and olivine from the “FS” parental melt.

IMPLICATIONS FOR MODERN ERUPTIONS

The evolution of the “FS” parental melt upon decompression, cooling, degassing and extensive (50%–60%) crystallization of olivine and clinopyroxene (with the plagioclase liquidus depressed by high H₂O contents) can hypothetically account for the compositions of modern lavas at Mt. Etna. The latter show the compositional trends in major and trace elements that can be tracked back to the “FS” compositions (Fig. 2), the geochemical signature consistent with their derivation from the high-magnesian, “FS”-type magma (Fig. 1). Water and CO₂ contents in the melt inclusions of modern eruptions (Del Carlo and Pompilio, 2004; Métrich et al., 2004; Métrich and Rutherford, 1998; Spilliaert et al., 2006) also point to the volatile-enriched character of parental magmas (Fig. 3), however linking the “FS” magmas and the present day magmas requires a more comprehensive interpretation of the behavior of volatile elements. Quantitative relationships between the “FS” and modern magmas are hampered by current lack of relevant experimental data to model the olivine-clinopyroxene cotectic, and undefined extents of primitive magma replenishment, interaction with earlier cumulates, mixing with evolved magmas in the conduit and degassing.

In historic times, and most evident recently, Mt. Etna showed prominent geochemical changes in the erupted products (Armienti et al., 2004; Clocchiatti et al., 1988; Tanguy et al., 1997; Tonarini et al., 2001) and dramatically increased volcanic activity, seismicity and ground deformation (Branca and Del Carlo, 2005; Patane et al., 2003). Uncommon elsewhere in the world, frequent explosive eruptions of basaltic

magmas in the latest volcanic cycles of Mt. Etna are coupled with the largest (for a single volcano) emissions of volcanic gases CO₂ and SO₂ (Allard, 1997; Allard et al., 1991; D'Alessandro et al., 1997). Another unusual aspect of the Mt. Etna magmatic system is the vertical extent (up to 18 km) of the magma storage region and related mafic cumulates advocated by geophysical and geochemical data (Allard, 1997; Chiarabba et al., 2000; De Gori et al., 2005; Laigle et al., 2000; Tanguy et al., 1997). A prerequisite to all features that make present-day Mt. Etna an extraordinary magmatic system is arrival of new primitive magma into the feeding system. Such magma, hypothesized in several works (Clocchiatti et al., 2004; Métrich et al., 2004; Patane et al., 2003; Spilliaert et al., 2006) but not sampled directly among recent volcanics, can be best approximated by an unevolved, mantle-derived "FS"-type magma, established in our study. The "FS"-type melt demonstrates properties critical for positive magma buoyancy; namely, originally high abundances of dissolved H₂O and CO₂ that were further increased by crystallization of olivine and clinopyroxene at depth. Subsequent decompression promoted early saturation of the melt in volatiles, vesiculation, and rapid ascent of the low-density, highly buoyant melt-bubble mixture. Thus, the explosive dynamics of modern Mt. Etna can be related to a deep supply of volatile-rich, high-Mg magmas. The recognition of such magmas in the plumbing system of Mt. Etna is also important in linking shallow-crust and surface processes to the evolution of the mantle source region. The "FS"-type melt provides compelling compositional evidence that Mt. Etna is currently supplied from a source, increasingly fluxed by melts and fluids, in a continuously changing tectonic framework.

ACKNOWLEDGMENTS

183 This paper has greatly benefited from comments by Malcolm Rutherford, Paul
184 Wallace and Dawnika L. Blatter. We thank the editor Andrew Barth for improving
185 the manuscript. This research was supported by a Research Fellowship and Discovery
186 grant (ARC, Australia) and F.W. Bessel Award (A. von Humboldt Foundation,
187 Germany) to V. Kamenetsky and an INGV-DPC grant to M. Pompilio.

188 **REFERENCES CITED**

- 189 Allard, P., 1997, Endogenous magma degassing and storage at Mount Etna: Geophysical
190 Research Letters, v. 24, p. 2219–2222, doi: 10.1029/97GL02101.
- 191 Allard, P., Carbonnelle, J., Dajlevic, D., Lebronec, J., Morel, P., Robe, M.C., Maurenas,
192 J.M., Faivrepierrret, R., Martin, D., Sabroux, J.C., and Zettwoog, P., 1991, Eruptive
193 and diffuse emissions of CO₂ from Mount Etna: Nature, v. 351, p. 387–391, doi:
194 10.1038/351387a0.
- 195 Andronico, D., Branca, S., Calvari, S., Burton, M., Caltabiano, T., Corsaro, R.A., Del
196 Carlo, P., Garfi, G., Lodato, L., Miraglia, L., Mure, F., Neri, M., Pecora, E.,
197 Pompilio, M., Salerno, G., and Spampinato, L., 2005, A multi-disciplinary study of
198 the 2002–03 Etna eruption: insights into a complex plumbing system: Bulletin of
199 Volcanology, v. 67, p. 314–330, doi: 10.1007/s00445-004-0372-8.
- 200 Armienti, P., Tonarini, S., D’Orazio, M., and Innocenti, F., 2004, Genesis and evolution
201 of Mt. Etna alkaline lavas: petrological and Sr-Nd-B isotope constraints: Periodico di
202 Mineralogia, v. 73, p. 29–52.
- 203 Branca, S., and Del Carlo, P., 2005, Types of eruptions of Etna volcano AD 1670–2003:
204 implications for short-term eruptive behaviour: Bulletin of Volcanology, v. 67,
205 p. 732–742, doi: 10.1007/s00445-005-0412-z.

- 206 Chiarabba, C., Amato, A., Boschi, E., and Barberi, F., 2000, Recent seismicity and
207 tomographic modeling of the Mount Etna plumbing system: *Journal of Geophysical*
208 *Research*, v. 105, p. 10923–10938, doi: 10.1029/1999JB900427.
- 209 Clocchiatti, R., Condomines, M., Guenot, N., and Tanguy, J.C., 2004, Magma changes at
210 Mount Etna: the 2001 and 2002–2003 eruptions: *Earth and Planetary Science*
211 *Letters*, v. 226, p. 397–414, doi: 10.1016/j.epsl.2004.07.039.
- 212 Clocchiatti, R., Joron, J.L., and Treuil, M., 1988, The role of selective alkali
213 contamination in the evolution of recent historic lavas of Mt Etna: *Journal of*
214 *Volcanology and Geothermal Research*, v. 34, p. 241–249, doi: 10.1016/0377-
215 0273(88)90036-4.
- 216 Coltelli, M., Del Carlo, P., Pompilio, M., and Vezzoli, L., 2005, Explosive eruption of a
217 picrite: The 3930 BP subplinian eruption of Etna volcano (Italy): *Geophysical*
218 *Research Letters*, v. 32, p. L23307, doi: 10.1029/2005GL024271.
- 219 Coltelli, M., Del Carlo, P., and Vezzoli, L., 2000, Stratigraphic constraints for explosive
220 activity in the past 100 ka at Etna Volcano, Italy: *International Journal of Earth*
221 *Sciences*, v. 89, p. 665–677, doi: 10.1007/s005310000117.
- 222 Corsaro, R.A., Miraglia, L., and Pompilio, M., 2006, Petrologic evidence of a complex
223 plumbing system feeding the July-August 2001 eruption of Mt. Etna, Sicily, Italy:
224 *Bulletin of Volcanology*, in press, doi: 10.1007/s00445-006-0083-4.
- 225 Corsaro, R.A., and Pompilio, M., 2004, Dynamics of magma at Mount Etna, *in*
226 Bonaccorso, A., et al., eds., *Mt Etna: Volcano Laboratory*, Volume 143: Washington,
227 DC, American Geophysical Union, p. 91–110.

- 228 D'Alessandro, W., Giammanco, S., Parello, F., and Valenza, M., 1997, CO₂ output and
229 $\delta^{13}\text{C}(\text{CO}_2)$ from Mount Etna as indicators of degassing of shallow asthenosphere:
230 Bulletin of Volcanology, v. 58, p. 455–458, doi: 10.1007/s004450050154.
- 231 De Gori, P., Chiarabba, C., and Patane, D., 2005, *Qp* structure of Mount Etna:
232 Constraints for the physics of the plumbing system: Journal of Geophysical
233 Research, v. 110, p. B05303, doi: 10.1029/2003JB002875.
- 234 Del Carlo, P., and Pompilio, M., 2004, The relationship between volatile content and the
235 eruptive style of basaltic magma: the Etna case: Annals of Geophysics, v. 47,
236 p. 1423–1432.
- 237 Doglioni, C., Innocenti, F., and Mariotti, G., 2001, Why Mt Etna?: Terra Nova, v. 13,
238 p. 25–31, doi: 10.1046/j.1365-3121.2001.00301.x.
- 239 Grove, T.L., Kinzler, R.J., and Bryan, W.B., 1992, Fractionation of mid-ocean ridge
240 basalt (MORB), in Phipps Morgan, J., Blackman, D.K., and Sinton, J., eds., Mantle
241 Flow and Melt Generation at Mid-Ocean Ridges, AGU Geophysical Monograph,
242 v. 71: Washington, D.C., p. 281–310.
- 243 Gueguen, E., Doglioni, C., and Fernandez, M., 1998, On the post-25 Ma geodynamic
244 evolution of the western Mediterranean: Tectonophysics, v. 298, p. 259–269, doi:
245 10.1016/S0040-1951(98)00189-9.
- 246 Gvirtzman, Z., and Nur, A., 1999, The formation of Mount Etna as the consequence of
247 slab rollback: Nature, v. 401, p. 782–785, doi: 10.1038/44555.
- 248 Kamenetsky, V., and Clocchiatti, R., 1996, Primitive magmatism of Mt Etna: insights
249 from mineralogy and melt inclusions: Earth and Planetary Science Letters, v. 142,
250 p. 553–572, doi: 10.1016/0012-821X(96)00115-X.

- 251 Kamenetsky, V.S., Crawford, A.J., and Meffre, S., 2001, Factors controlling chemistry of
252 magmatic spinel: an empirical study of associated olivine, Cr-spinel and melt
253 inclusions from primitive rocks: *Journal of Petrology*, v. 42, p. 655–671, doi:
254 10.1093/petrology/42.4.655.
- 255 Laigle, M., Hirn, A., Sapin, M., Lepine, J.C., Diaz, J., Gallart, J., and Nicolich, R., 2000,
256 Mount Etna dense array local earthquake P and S tomography and implications for
257 volcanic plumbing: *Journal of Geophysical Research*, v. 105, p. 21633–21646, doi:
258 10.1029/2000JB900190.
- 259 Marty, B., Trull, T., Lussiez, P., Basile, I., and Tanguy, J.C., 1994, He, Ar, O, Sr and Nd
260 isotope constraints on the origin and evolution of Mount Etna magmatism: *Earth and*
261 *Planetary Science Letters*, v. 126, p. 23–39, doi: 10.1016/0012-821X(94)90240-2.
- 262 Métrich, N., Allard, P., Spilliaert, N., Andronico, D., and Burton, M., 2004, 2001 flank
263 eruption of the alkali- and volatile-rich primitive basalt responsible for Mount Etna's
264 evolution in the last three decades: *Earth and Planetary Science Letters*, v. 228, p. 1–
265 17, doi: 10.1016/j.epsl.2004.09.036.
- 266 Métrich, N., and Rutherford, M.J., 1998, Low pressure crystallization paths of H₂O-
267 saturated basaltic-hawaiitic melts from Mt Etna: Implications for open-system
268 degassing of basaltic volcanoes: *Geochimica et Cosmochimica Acta*, v. 62, p. 1195–
269 1205, doi: 10.1016/S0016-7037(98)00048-9.
- 270 Newman, S., and Lowenstern, J.B., 2002, VolatileCalc: a silicate melt-H₂O-CO₂ solution
271 model written in Visual Basic for excel: *Computers & Geosciences*, v. 28, p. 597–
272 604, doi: 10.1016/S0098-3004(01)00081-4.

- 273 Patane, D., De Gori, P., Chiarabba, C., and Bonaccorso, A., 2003, Magma ascent and the
274 pressurization of Mount Etna's volcanic system: *Science*, v. 299, p. 2061–2063, doi:
275 10.1126/science.1080653.
- 276 Schiano, P., Clocchiatti, R., Ottolini, L., and Busa, T., 2001, Transition of Mount Etna
277 lavas from a mantle-plume to an island-arc magmatic source: *Nature*, v. 412, p. 900–
278 904, doi: 10.1038/35091056.
- 279 Spilliaert, N., Allard, P., Métrich, N., and Sobolev, A.V., 2006, Melt inclusion record of
280 the conditions of ascent, degassing and extrusion of volatile-rich alkali basalt during
281 the powerful 2002 flank eruption of Mount Etna (Italy): *Journal of Geophysical*
282 *Research*, v. 111, p. B04203, doi: 10.1029/2005JB003934.
- 283 Sun, S.-S., and McDonough, W.F., 1989, Chemical and isotopic systematics of oceanic
284 basalts: implications for mantle composition and processes, *in* Saunders, A.D., and
285 Norry, M.J., eds., *Magmatism in the Ocean Basins*, London, Geological Society
286 Special Publication, v. 42, p. 313–345.
- 287 Tanguy, J.C., Condomines, M., and Kieffer, G., 1997, Evolution of the Mount Etna
288 magma: Constraints on the present feeding system and eruptive mechanism: *Journal*
289 *of Volcanology and Geothermal Research*, v. 75, p. 221–250, doi: 10.1016/S0377-
290 0273(96)00065-0.
- 291 Tonarini, S., Armienti, P., D'Orazio, M., and Innocenti, F., 2001, Subduction-like fluids
292 in the genesis of Mt. Etna magmas: evidence from boron isotopes and fluid mobile
293 elements: *Earth and Planetary Science Letters*, v. 192, p. 471–483, doi:
294 10.1016/S0012-821X(01)00487-3.

Wallace, P.J., 2005, Volatiles in subduction zone magmas: concentrations and fluxes based on melt inclusion and volcanic gas data: *Journal of Volcanology and Geothermal Research*, v. 140, p. 217-240; doi: 10.1016/j.jvolgeores.2004.07.023.

FIGURE CAPTIONS

Figure 1. Primitive mantle normalized (after Sun and McDonough, 1989) trace element compositions of “FS” rocks (open circles, average of 6 samples), melt inclusions hosted in “FS” olivine phenocrysts (closed circles, average of 23 melt inclusions; the bars show one standard deviation), and Mount Etna lavas erupted in 2001–2002 (diamonds, average of 61 samples; after Andronico et al., 2005; Corsaro et al., 2006).

Figure 2. Comparison between the compositions of the “FS” parental magmas (exemplified by recalculated melt inclusions), “FS” whole rocks and modern lavas of Mount Etna (Andronico et al., 2005; Corsaro et al., 2006). Note significant variations in trace elements among individual melt inclusions. Symbols as in Figure 1.

Figure 3. Relationships between H₂O and CO₂ contents in melt inclusions from “FS” olivine (circles), phenocrysts in recent Mount Etna lavas (diamonds; after Métrich et al., 2004; Spilliaert et al., 2006) and arc basalts (dotted outline; after Wallace, 2005). Note that adequate comparison of data from different sources requires normalization to the same degree of fractionation. Isobars are calculated at 1220 °C and 47.5 wt% SiO₂ in melt using VOLATILECALC (Newman and Lowenstern, 2002). Inset shows typical H₂O- and CO₂-rich silicate melt inclusion with a CO₂-vapor bubble in “FS” olivine. The bubble’s walls are coated by tiny Ca-Mg-bearing carbonates (determined by laser Raman

318 spectroscopy, GeoForschungsZentrum Potsdam, Germany), thus the initial CO₂ content
319 was even higher than measured.

320

321 ¹GSA Data Repository item 2007xxx, Analytical and calculation methods, Tables DR1,
322 DR2 and DR3, is available online at www.geosociety.org/pubs/ft2007.htm, or on request
323 from editing@geosociety.org or Documents Secretary, GSA, P.O. Box 9140, Boulder,
324 CO 80301, USA.